



# Astronomical League Southern Sky Telescopic Club



Many of us in the Northern hemisphere dream of a vacation to the Southern hemisphere and observing under the legendary Southern Skies. We may be lucky enough to go south once or twice in a lifetime. When we get that chance, and have access to a telescope, we want a list of interesting objects at hand. A reminder of the objects we don't want to miss. That's the idea behind this selection of Southern Sky Showpieces.

There are 100 objects total on the attached lists. These are broken down into 20 "far-south" objects (-65 to -90), 40 "mid-south" objects (-48 to -65), and 40 "others" (-11 to -47). We tried to include the generally accepted "best" objects in the far south areas in the sky. The "other" objects north of -48 degrees are a mixed group. These objects include some old favorite Messier objects as well as more unfamiliar objects for northern observers.

## Observing the Objects

All the objects on this list will be visible in a small telescope in good dark skies. Although many are visible in large binoculars, a telescope is very highly recommended for this project. Those wishing to observe with binoculars should pursue the Astronomical League Southern Sky Binocular list rather than this list. Particularly with the planetary nebula, higher magnifications are needed than can be provided by binoculars. The galaxies on the list will also benefit greatly from the added aperture of a telescope. All of these objects can be seen in a 5 inch (125mm) or larger telescope. Certainly more than half can be seen in an 80mm scope under dark skies.

Many of the objects in this observing program are also in the Astronomical League's Southern Skies Binocular, Caldwell, and Messier observing programs. There are even a few bright Arp objects. Any observer intending to use the same set of observations for multiple programs must keep in mind the varying criteria. The Messier and Caldwell programs, for example, do not allow computer controlled (go-to) telescopes.

You will need a star chart to find the objects. If you have a computer charting program, you might wish to print out charts to help you find the objects. Plan ahead to maximize your observing time, you will need to know what will be visible at your site. At some times of the year, such as late September, it may be difficult to view the required number of objects without careful planning. Some objects may not be listed in your atlas or computer program at all, or an alternate name may be used. In that case, look at your charts for the other objects in the group and check at the given coordinates.

## To earn a Southern Sky Showpiece Award

- You must be a member of the Astronomical League,
- You must locate, observe, and log at least 50 objects. Of these 50, at least 30 must be south of -48 degrees, and at least 10 of those 30 must be south of -65 degrees.
- "Go-to" telescopes are allowed.
- Keep a log recording your observations including:
  - Object's name/number
  - Date and time
  - Your observing location
  - Seeing and/or transparency
  - Approximate altitude of the object above the horizon
  - Size of telescope used and magnification(s)
  - Any filters used
  - Description of object

If you belong to a club that is affiliated with the Astronomical League, one of your club officers will verify your observations and notify the AL. If you do not belong to a club, but join the Astronomical League as a member at large, you can send your observations directly to the Coordinator at the address given in the current issue of the Reflector or on the League web page. Send only copies of your observations in case they get lost in the mail, and remember to mention that you are a League member at large.

In return, you will receive a certificate, and a lapel pin celebrating your accomplishment.

## Your Observing Log

Seeing-- is an indicator of how steady the atmosphere is. There are many excellent ways to evaluate seeing.

For example- a scale of 1-5 with 1 being a perfectly steady image and 5 being violent turbulence.

Transparency- this is a measure of how faint you can see. This can be affected by humidity, dust, natural sky glow, moonlight, pollution, and observing altitude. Estimating the magnitude of the faintest star you can see is a simple measure of transparency. Whatever scale you use, try to be consistent.

Object altitude above the horizon:

The simplest way to estimate an object's approximate altitude above the horizon is the "hand method": Hold your hand out at full arm's length-the distance between the tips of your extended thumb and extended little finger is about 20 degrees. Your closed fist is about 10 degrees, and three fingers is about five. If you didn't estimate the altitude above the horizon when you made your observations, you can 'reconstruct' the information from the date, time, and location of the observations.

## About the Objects:

The choice of objects for any observing program is highly subjective. You will no doubt find some great objects that were overlooked, and some duds that were included. Beyond a few obvious choices (such as 47 Tucanae and the Eta Carina nebula) various observers will have different favorites. For an observing program such as this, we also wanted to have objects available at any time of year for a dedicated observer to complete the program in one trip South. And, we wanted the list of objects to include an assortment of types of objects. The list is heavy on globular clusters, as these wondrous balls of stars are concentrated around the center of our galaxy-- and the center of our galaxy is south. The list also includes several "objects" that are really groups or pairs of objects that are visible in one low-power eyepiece field of view. For purposes of the observing award, you only need to see one of the objects in the group. Most of the time, you should be able to see them all.

Far South: -65 to -90 degrees; The Magellanic Clouds and South

This group of objects includes 47 Tucanae, the second brightest globular in the sky, the Small Magellanic Cloud (SMC), and the Large Magellanic Cloud (LMC). This is the territory for observing objects in other galaxies!

The SMC is relatively poor in easily visible star clusters and nebula. By contrast, the LMC is a wonderfully rich area to explore. It would be easy to spend weeks viewing the clusters and nebula in the LMC. By far the most notable nebulosity in the LMC is the Tarantula Nebula. The list includes it, and some interesting fields in the LMC. You will find that it is difficult to find a field with just one or two patches of stars or nebula. Enjoy!

Mid-South: -48 to -65 degrees; The Southern Milky Way Band

This area is between Omega Centauri and the Magellanic Clouds. It includes a swath of the Southern Milky way including Eta Carina, the Coal Sack and the Jewel Box cluster. This is an area very rich in open clusters of all sizes. If you are lucky enough to observe in truly dark transparent skies, look for the dark nebula tracking through the Milky Way and bordering many of the open clusters. Three double stars and two dark nebula are included in the group for a change of pace.

Proxima Centauri, the nearest star to our sun, is a very faint 11th magnitude companion of Alpha Centauri. It is about 2 degrees from Alpha, and impossible to identify with certainty without a detailed finder chart. For that reason, it is not included on this list.

Keyhole Nebula-- the Keyhole nebula is a small, curved, dark dust lane crossing the brightest portion of the Eta Carina nebula. It is located west of the star Eta Carina. The keyhole shape has become less apparent over the years because of changes in Eta Carina and the associated nebulosity. A filter may make the dark nebula easier to spot by increasing the contrast between the bright and dark nebula.

The Other Group: -47 to -11 degrees:

This group includes Omega Centauri, Centaurus A, the Helix and a number of Messier objects.

Why include objects that can be seen from mid-north latitudes at all? Atmosphere! If you observe an object at just 30 degrees above the horizon, you are looking through twice the distorting layer of atmosphere than if the object is directly overhead. That additional layer of atmosphere causes a significant loss in detail. These familiar objects are on the list in the hope that you will take the time to look at them when they are at their best-high above the horizon. Observe them when they are 45 degrees or more above the horizon and you will be rewarded with a new appreciation of your old favorites.

## Dedication:

This observing program is dedicated to the memory of the late Ken Wilcox- who had the courage to start the Southern Skies Star Party in Bolivia. Great thanks are also due Vic and Jen Winter, who have ably continued and expanded on Ken's vision.

Kathy Machin, Astronomical Society of Kansas City; 4845 N Smalley, Kansas City MO 64119; [kmachin@kc.rr.com](mailto:kmachin@kc.rr.com)

The Far South Group [-65 and South]

Observe and Log at least 10 of these 20 objects

	Name	Con	RA	DEC	Mag	Size	Type	Notes
[ ]	NGC 104	Tuc	00h 24.1 m	-72° 05'	4	50'	GC	47 Tucanae
[ ]	NGC 292	Tuc	00h 52.8 m	-72° 50'	2.7	210x140'	Galaxy	The Small Magellanic Cloud (SMC)
[ ]	NGC 330	Tuc	00h 56.3 m	-72° 28'	9.6	2'	OC	In the SMC
[ ]	NGC 346	Tuc	00h 59.1 m	-72° 11'	10.3	14'	OC + Neb	In the SMC
[ ]	NGC 362	Tuc	01h 03.2 m	-70° 51'	6.8	13'	GC	
[ ]	NGC 371	Tuc	01h 03.5 m	-72° 05'		8'	OC	In the SMC
[ ]	NGC 419	Tuc	01h 08.3 m	-72° 53'	10.6	2.6'	GC	In the SMC
[ ]	NGC 1313	Ret	03h 18.2 m	-66° 30'	9.2	9.1x6.9'	Galaxy	
[ ]	NGC 1763 group	Dor	04h 56.8 m	-66° 24'		25.0'	Br Neb	In LMC, with NGC 1760, 61, 69, 73
[ ]	NGC 1850 group	Dor	05h 08.7 m	-68° 46'	9.3	3'	OC	In the LMC, group with NGC 1854,58,63
[ ]	NGC 1955 group	Dor	05h 26.1 m	-67° 28'	9		OC + Neb	In the LMC, group with NGC 1968,74
[ ]	NGC 1962 group	Dor	05h 26.5 m	-68° 46'	8		OC + Neb	In the LMC, group with NGC 1965,66,70
[ ]	NGC 2070	Dor	05h 38.7 m	-69° 06'	8.2	40'	OC + Neb	In the LMC, The Tarantula
[ ]	NGC 2074 group	Dor	05h 38.9 m	-69° 28'			OC + Neb	In the LMC, group with NGC 2081
[ ]	NGC 3195	Cha	10h 09.4 m	-80° 52'	11.5	0.6'	PN	
[ ]	NGC 4372	Mus	12h 25.8 m	-72° 40'	7.2	5'	GC	
[ ]	NGC 4833	Mus	12h 59.6 m	-70° 53'	8.4	13.5'	GC	
[ ]	NGC 5189	Mus	13h 33.5 m	-65° 59'	9.5	3.1x1.2'	PN	
[ ]	NGC 6101	Aps	16h 25.8 m	-72° 12'	9.2	5'	GC	
[ ]	NGC 6362	Ara	17h 31.9 m	-67° 03'	8.1	14'	GC	

Southern Milky Way Band [-48° to -64°]

Observe at least 30 total from this group plus the Far South Group.

	Name	Con	RA	DEC	Mag	Size	Type	Notes
[ ]	Beta Tuc	Tuc	00h 31.5 m	-62° 58'			Double Star	4.4, 4.5 27"; 10' 5.1"
[ ]	NGC 1261	Hor	03h 12.3 m	-55° 13'	8.3	7'	GC	
[ ]	NGC 1549/1553	Dor	04h 15.7 m	-55° 35'	10.7	4.9x4'	Galaxy pair	
[ ]	NGC 1566	Dor	04h 20.0 m	-54 56'	10.3	8.3x 6.5'	galaxy	
[ ]	NGC 2516	Car	07h 58.3 m	-60° 51'	3.8	30'	OC	
[ ]	NGC 2547	Vel	08h 10.2 m	-49° 14'	4.7	20'	OC	
[ ]	IC 2395	Vel	08h 42.5 m	-48° 07'	4.6	8'	OC	
[ ]	NGC 2669	Vel	08h 46.3 m	-52° 52'	6.1	12'	OC	
[ ]	NGC 2808	Car	09h 12.0 m	-64° 52'	6.2	13'	GC	
[ ]	NGC 3114	Car	10h 02.7 m	-60° 06'	4.2	35'	OC	
[ ]	NGC 3247	Car	10h 25.9 m	-57° 56'	7.6	6'	OC + Neb	
[ ]	NGC 3293	Car	10h 35.8 m	-58° 14'	4.7	40.0'	OC + Neb	
[ ]	NGC 3324	Car	10h 37.3 m	-58° 38'	6.7	16.0'	OC + Neb	
[ ]	NGC 3372	Car	10h 43.8 m	-59° 52'		120.0'	OC + Neb	Eta Carina Nebula
[ ]	IC 2602	Car	10h 43.9 m	-64° 24'	1.6	50'	OC	Southern Pleiades
[ ]	Keyhole Neb	Car	10h 44.3 m	-59° 39'			Dk Nebula	Dk neb near brightest part of NGC 3372
[ ]	NGC 3532	Car	11h 05.6 m	-58° 44'	3	55.0'	OC	
[ ]	NGC 3572	Car	11h 10.4 m	-60° 15'	6.6	20.0'	OC + Neb	
[ ]	NGC 3766	Cen	11h 36.1 m	-61° 37'	5.3	12.0'	OC	
[ ]	Cr 249	Cen	11h 38.3 m	-63° 22'	4.5	75.0'	OC + Neb	Lambda Cen OC + Neb IC2944, 2948
[ ]	NGC 3918	Cen	11h 50.3 m	-57° 11'	8.4	0.3'	PN	Blue Planetary

[ ]	Acrux	Cru	12 h 26.6m	-63° 06'			Double Star	1.3, 1.7 4.2"; 4.9 90"
[ ]	NGC 4609	Cru	12h 42.3 m	-62° 59'	6.5	5'	OC	
[ ]	Coalsack	Cru	12h 53 m	-63°		400'	Dk Nebula	Very large Dark Nebula in Crux
[ ]	NGC 4755	Cru	12h 53.6 m	-60° 22'	4.2	10'	OC	The Jewel Box, Kappa Crucis Cluster
[ ]	NGC 4945	Cen	13h 05.4 m	-49° 28'	9.3	20x3.8'	Galaxy	
[ ]	NGC 5286	Cen	13h 46.4 m	-51° 22'	7.4	11'	GC	
[ ]	NGC 5281	Cen	13h 46.6 m	-62° 55'	5.9	5.0'	OC	
[ ]	NGC 5316	Cen	13h 54.0 m	-61° 52'	6	13'	OC	
[ ]	Alpha Centauri	Cen	14h 39.7 m	-60° 50'	0		Double star	0.0, 1.2 mag; separation 18.1"
[ ]	NGC 5823	Cir	15h 05.5 m	-55° 36'	7.9	10.0'	OC	
[ ]	NGC 5927	Lup	15h 28.0 m	-50° 40'	8	5.7'	GC	
[ ]	NGC 5946	Nor	15h 35.5 m	-50° 40'	8.4	3'	GC	
[ ]	NGC 6025	TrA	16h 03.3 m	-60° 25'	5.1	12'	OC	
[ ]	NGC 6067	Nor	16h 13.2 m	-54° 13'	5.6	12'	OC	
[ ]	NGC 6087	Nor	16h 18.9 m	-57° 54'	5.4	12'	OC	
[ ]	NGC 6352	Ara	17h 25.5 m	-48° 25'	7.8	9'	GC	
[ ]	NGC 6397	Ara	17h 40.7 m	-53° 40'	5.3	30'	GC	
[ ]	NGC 6744	Pav	19h 09.8 m	-63° 51'	9.1	20x12'	Galaxy	
[ ]	NGC 6752	Pav	19h 10.9 m	-59° 59'	5.3	28'	GC	

### The Rest of The Southern Skies List

## Observe a maximum of 20 from this group

	Name	Con	RA	DEC	Mag	Size	Type	Notes
[ ]	NGC 0055	Scl	00h 15.1 m	-39° 12'	8.4	32x6'	Galaxy	
[ ]	NGC 0253	Scl	00h 47.6 m	-25° 17'	8	27x6'	Galaxy	The Sculptor Galaxy
[ ]	NGC 0288	Scl	00h 52.8 m	-26° 35'	8.1	13'	GC	
[ ]	NGC 1097	For	02h 46.3 m	-30° 16'	10.2	9.3 x 6.2'	Galaxy	
[ ]	NGC 1291	Eri	03h 17.3 m	-41° 07'	9.4	9.8x8.1'	Galaxy	
[ ]	NGC 1316	For	03h 22.6 m	-37° 12'	9.4	11x7'	Galaxy	
[ ]	NGC 1360	For	03h 33.3 m	-25° 52'	9.6	6.4'	PN	
[ ]	NGC 1365	For	03h 33.6 m	-36° 08'	10.3	11.3x6.2'	Galaxy	
[ ]	NGC 1399 cluster	For	03h 38.5 m	-35° 27'	9.9	3.2x3.1'	Galaxy	Fornax Gal. with NGC's 1380, 1404
[ ]	NGC 1851	Col	05h 14.1 m	-40° 03'	7.01	11.0'	GC	
[ ]	NGC 2439	Pup	07h 40.8 m	-31° 41'	6.9	10'	OC	
[ ]	NGC 2447	Pup	07h 44.6 m	-23 52'	6.2	22'	OC	M 93
[ ]	NGC 2527	Pup	08h 04.9 m	-28 09'	6.5	22'	OC	
[ ]	NGC 3132	Vel	10h 07.0 m	-40° 26'	8.2	1.3x0.8'	PN	Eight-burst planetary
[ ]	NGC 3201	Vel	10h 17.6 m	-46° 25'	6.9	21'	GC	
[ ]	NGC 3242	Hya	10h 24.8 m	-18° 38'	8.6	.4	PN	Ghost of Jupiter
[ ]	NGC 3621	Hya	11h 18.3 m	-32 48'	9.8	13.3x6.1'	Galaxy	
[ ]	NGC 4361	Crv	12h 24.5 m	-18 47'	10	1.7'	PN	
[ ]	NGC 4594	Vir	12h 40.0 m	-11° 37'	9	8.8x3.5'	Galaxy	M 104 Sombrero
[ ]	NGC 5128	Cen	13h 25.5 m	-43° 01'	7.8	25x20'	Galaxy	Centaurus A
[ ]	NGC 5139	Cen	13h 26.8 m	-47° 29'	3.9	53'	GC	Omega Centauri
[ ]	NGC 5236	Hya	13h 37.0 m	-29° 52'	8.2	12.8x11.4'	Galaxy	M 83
[ ]	NGC 5986	Lup	15h 46.1 m	-37° 47'	7.6	9'	GC	
[ ]	NGC 6121	Sco	16h 23.6 m	-26° 32'	5.4	30'	GC	M 4
[ ]	NGC 6231	Sco	16h 54.2 m	-41° 49'	2.6	14'	OC + Neb	

[ ]	NGC 6250	Ara	16h 58.0 m	-45° 56'	5.9	7'	OC	
[ ]	NGC 6266	Oph	17h 01.2 m	-30° 07'	6.4	11'	GC	M 62
[ ]	NGC 6281	Sco	17h 04.8 m	-37° 54'	5.4	8'	OC + Neb	
[ ]	NGC 6388	Sco	17h 36.3 m	-44° 44'	6.8	6'	GC	
[ ]	NGC 6405	Sco	17h 40.3 m	-32° 16'	4.2	14'	OC	M 6
[ ]	NGC 6475	Sco	17h 53.9 m	-34° 49'	3.3	80.0'	OC	M 7
[ ]	NGC 6514	Sgr	18h 02.3 m	-23° 01'	6	17x20'	OC + Neb	M 20
[ ]	B 86	Sgr	18h 02.9 m	-27° 52'		5'	Dk Neb	Near NGC 6520
[ ]	NGC 6523	Sgr	18h 03.8 m	-24° 23'	6	50x40'	Br Neb	M 8
[ ]	NGC 6618	Sgr	18h 20.8 m	-16° 11'	6	46.0'	OC + Neb	M 17
[ ]	NGC 6656	Sgr	18h 36.4 m	-23° 54'	5.2	33'	GC	M 22
[ ]	NGC 6723	Sgr	18h 59.6 m	-36° 38'	6.8	13'	GC	Neb NGC 6726-27,29 1/2 degree SE
[ ]	NGC 6809	Sgr	19h 40.0 m	-30° 58'	6.3	19'	GC	M 55
[ ]	NGC 7293	Aqr	22h 29.6 m	-20° 48'	7.5	16'	PN	Helix Nebula
[ ]	NGC 7793	Scl	23h 57.8 m	-32 35'	9.6	10.0x6.5	Galaxy	